

Deworming and Vaccination Schedules for Wild Horses and Burros at Off-Range Corrals**Table 1: Deworming Schedule for Wild Horses and Burros at Off-Range Corrals**

Treatment	Adult Older Than 1 Year		Foals (Born in the Current Foaling Year)	
	Deworming ¹	Primary dose	Every 180 days	Primary dose when first vaccinated, repeated at time of second booster

1. Ivermectin liquid will be the primary medication used for routine deworming. Moxidectin or another type of dewormer may also be used in specific cases if recommended by an attending veterinarian.

Table 2: Vaccination Schedule for Wild Horses and Burros at Off-Range Corrals

Vaccines ²	Adult (Older than 1 year)		Foals (Born in current foaling year) (Primary given at 4-6 months of age)	
	Primary and First Booster	Maintenance Booster	Primary and First Booster	Second Foal Booster and Maintenance Booster
Influenza (Equine flu) Rhinopneumonitis (Equine herpes) Strangles (Strep. equi)	Primary + 1 st booster (3-4 weeks apart)	Every 180 days	Primary + 1 st booster (3-4 weeks apart)	2 nd Booster 30-90 days after 1 st booster After that, every 180 days
Eastern & Western Equine Encephalomyelitis Tetanus West Nile Virus	Primary + 1 st booster (3-4 weeks apart)	Once a year	Primary + 1 st booster (3-4 weeks apart)	2 nd Booster 30-90 days after 1 st booster After that once a year
Rabies	Primary	Once a year	Primary + 1 st Booster 30-90 days apart	After that once a year

2. In the case of limited vaccine availability due to manufacturing or supply backorders, BLM personnel will consult with the WHB Program's veterinary advisor to adjust this vaccination schedule as needed.